NUTRITIONAL STRATEGIES FOR ENHANCED RECOVERY AND PERFORMANCE IN CROSS-RUNNING

Khaydarov Sayfulloh

Received: 22.10.2025
Revised: 23.10.2025
Accepted: 24.10.2025

Keywords:
cross-country running, nutrition, recovery, protein intake 2.2 g/kg, carbohydrates, antioxidants, omega-3, DOMS

Abstract: This thesis explores the impact of nutritional strategies on post-exercise recovery and performance optimization among cross-runners. Proper nutrition is fundamental not only for maintaining energy balance during endurance training but also for facilitating efficient muscle regeneration, immune system resilience, and overall physiological adaptation following competition. The study underscores the significance of the *"nutritional window of opportunity"* — a critical period immediately after exercise when timely consumption of carbohydrates, proteins, and antioxidants can markedly enhance recovery efficiency.

An experimental survey involving 67 cross-runners demonstrated that post-race nutrition exerts a decisive influence on reducing fatigue, alleviating muscle soreness, and accelerating recovery. Among participants, 53.7% consumed protein-rich foods, 25.4% relied on carbohydrate-based meals, 14.9% used recovery drinks, and 6% preferred complex beverages containing a combination of protein, carbohydrates, and electrolytes. Athletes who consumed approximately 2.2 grams of protein per kilogram of body weight per day exhibited significantly faster recovery and reduced muscle damage, particularly when protein and carbohydrates were ingested together within two hours post-exercise.

Furthermore, diets enriched with calcium, vitamin D, omega-3 fatty acids, and natural antioxidants were associated with improved muscle repair, reduced inflammation, enhanced immune function, and lower risk of overuse injuries. These findings highlight that strategic nutritional planning, with special emphasis on adequate protein intake (~2.2 g/kg/day) and nutrient timing, constitutes an essential element of modern endurance training. Implementing these evidence-based nutritional approaches can substantially enhance both recovery efficiency and sustained athletic performance in cross-running athletes.

Introduction. Cross- running is one of the most physically demanding endurance sports, requiring a high level of aerobic capacity, muscular endurance, and metabolic efficiency. During intense and prolonged exercise, glycogen stores are depleted, muscle fibers experience microdamage, and oxidative stress increases — all of which negatively affect recovery and future performance.

Nutrition is therefore a fundamental factor in maintaining energy availability, accelerating recovery, and supporting the immune system of endurance athletes. According to Vitale and Getzin (2019), the "nutritional window of opportunity" — the first two hours after exercise — is when the body most effectively synthesizes glycogen and repairs muscle tissue. Moore et al. (2014) and Kerksick et al. (2017) further emphasize that sufficient protein and carbohydrate consumption during this period is crucial for muscle protein synthesis and recovery.

Recent studies also suggest that endurance athletes benefit from higher daily protein intake compared to non-athletes. Specifically, consuming about 2.0–2.2 g of protein per kilogram of body weight per day has been found to significantly enhance muscle recovery, prevent protein catabolism, and maintain lean body mass (Thomas et al., 2016).

Therefore, this study aims to assess the effects of post-exercise nutrition — particularly the role of protein intake at approximately 2.2 g/kg/day, in combination with carbohydrates, antioxidants, and micronutrients — on recovery efficiency among cross-country runners.\

Methodology:

Participants: The study involved **67 competitive cross-runners** aged 18–25 years, all of whom had at least two years of endurance training experience and no chronic health conditions. Written consent was obtained prior to participation.

Study Design

An **experimental descriptive design** was applied to analyze post-race nutrition effects over a four-week observation period. Participants were grouped by their preferred nutritional recovery method:

1. Protein-rich foods (53.7%)
2. Carbohydrate-focused meals (25.4%)
3. Recovery drinks (14.9%)
4. Complex beverages (6%)

Nutritional Intervention

Athletes were advised to consume nutrients within **two hours after training**. Each athlete's diet was individually adjusted to maintain an average daily protein intake of about **2.2 grams per kilogram of body weight**, divided evenly across meals every 3–4 hours. This ensured a consistent supply of amino acids to support muscle repair. They were also guided to:

- Maintain macronutrient distribution: 55–60% carbohydrates, 20–25% protein, 20% fat
- Include antioxidant-rich foods (berries, leafy greens, turmeric)
- Supplement omega-3 fatty acids and vitamin D

- Maintain hydration (12–14 cups water per day, with electrolytes when needed)

Data Collection

Data were collected through questionnaires, 5 km time trials (before and after intervention), observation of DOMS (delayed onset muscle soreness), and nutritional logs.

Results:

The findings confirmed that **protein intake of approximately 2.2 g/kg/day** significantly enhanced post-exercise recovery. Athletes following this intake pattern experienced less fatigue, reduced DOMS, and faster muscle regeneration compared to those consuming lower protein levels.



Overall, 81% of participants reported noticeable recovery improvements due to post-race nutrition. The distribution of nutritional approaches showed that:

- 53.7% consumed protein-rich meals
- 25.4% consumed carbohydrate-based foods
- 14.9% used recovery drinks
- 6% used complex beverages

Participants who consumed both **protein and carbohydrates within two hours post-training** recovered **34% faster**, while those who consistently met the **2.2 g/kg/day protein goal** showed **47% improvement** in muscle recovery and **30–35% reduction** in DOMS.

Antioxidant-rich and omega-3-supplemented diets further reduced inflammation (≈40%) and oxidative stress (≈50%). The inclusion of calcium and vitamin D supported bone strength and reduced injury risk by up to 15%

Discussion:

The results strongly support previous findings that emphasize the importance of protein timing and adequate intake for endurance athletes. Consuming approximately

2.2 g/kg/day of protein was associated with superior muscle recovery, reduced soreness, and greater training consistency — outcomes consistent with Kerksick et al. (2017) and Thomas et al. (2016).

Higher protein intake ensures optimal levels of leucine and other essential amino acids, stimulating muscle protein synthesis and reducing catabolic breakdown during recovery. Additionally, when protein is paired with carbohydrates shortly after exercise, glycogen resynthesis accelerates, further improving endurance performance.

The findings also align with prior studies on omega-3 and antioxidants (Howatson et al., 2010; Samaras et al., 2014), which highlight their anti-inflammatory properties and contribution to recovery efficiency. The synergy of macronutrients and micronutrients — particularly protein, carbohydrates, antioxidants, and vitamin D — forms the foundation of an effective recovery nutrition strategy.

In practical application, maintaining **daily protein intake of about 2.2 g/kg body weight,** spread evenly throughout the day, appears to be one of the most effective approaches to enhance recovery, prevent overtraining, and sustain high performance in cross-country runners.

Conclusion:

This study concludes that post-exercise nutrition plays a decisive role in the recovery, performance, and overall physiological adaptation of cross-country runners. The findings clearly demonstrate that athletes who adhered to a structured nutritional plan — particularly those consuming **approximately 2.2 grams of protein per kilogram of body weight per day** — experienced superior recovery outcomes compared to those with lower protein intake.

Adequate protein consumption, when paired with sufficient carbohydrate intake during the post-exercise "nutritional window," accelerated glycogen resynthesis, enhanced muscle repair, and minimized symptoms of delayed onset muscle soreness (DOMS). The study further confirmed that balanced macronutrient distribution, combined with antioxidant- and omega-3-rich foods, significantly reduces inflammation and oxidative stress, while maintaining muscle and bone integrity.

These results emphasize that the **quality, timing, and quantity** of nutrient intake are equally important in supporting endurance athletes' recovery and long-term performance. Therefore, integrating targeted nutritional strategies into daily training routines should be viewed as an essential part of modern sports science, not merely a supplementary component.

Recommendations:

1. **Protein Intake Optimization:** Cross-country runners should aim to consume **2.0–2.2 g of protein per kilogram of body weight per day**, distributed evenly across meals every 3–4 hours, to maximize muscle protein synthesis and enhance post-exercise recovery.
2. **Timing of Nutrient Intake:** Athletes should consume a combination of **high-quality proteins and**

carbohydrates within two hours after training or competition, as this period maximizes glycogen replenishment and tissue repair.

3. **Antioxidant and Omega-3 Supplementation:** Diets should include natural sources of antioxidants (such as berries, spinach, and turmeric) and omega-3 fatty acids (from fish oil or flaxseed), which reduce inflammation and support recovery efficiency.
4. **Micronutrient Support:** Ensuring sufficient intake of **calcium and vitamin D** is critical for bone strength and the prevention of stress injuries common in endurance athletes.
5. **Hydration and Electrolyte Balance:** Athletes should maintain consistent hydration (12–14 cups per day) and restore electrolyte balance, especially after long-distance sessions, to prevent muscle cramps and dehydration.
6. **Individualized Nutrition Plans:** Coaches and sports nutritionists should develop **personalized meal plans** that reflect each runner's training volume, metabolic rate, and recovery needs, ensuring optimal performance outcomes.
7. **Further Research:** Future studies should explore the long-term effects of different protein sources (plant-based vs. animal-based) and nutrient timing strategies on recovery, endurance performance, and injury prevention among endurance athletes.

REFERENCES:

Howatson, G., McHugh, M. P., Hill, J. A., Brouner, J., Jewell, A. P., Van Someren, K. A., Shave, R. E., & Howatson, S. A. (2010). *Influence of tart cherry juice on indices of recovery following marathon running. Scandinavian Journal of Medicine & Science in Sports, 20*(6), 843–852. https://doi.org/10.1111/j.1600-0838.2009.01005.x

Kerksick, C. M., Arent, S., Schoenfeld, B. J., Stout, J. R., Campbell, B., Wilborn, C. D., Taylor, L., Kalman, D., Smith-Ryan, A. E., Kreider, R. B., Willoughby, D., Arciero, P. J., VanDusseldorp, T. A., Ormsbee, M. J., Wildman, R., Greenwood, M., Ziegenfuss, T. N., Aragon, A. A., & Antonio, J. (2017). *International Society of Sports Nutrition position stand: Nutrient timing. Journal of the International Society of Sports Nutrition, 14*(1), 33–33. https://doi.org/10.1186/s12970-017-0189-4

Moore, D. R., Camera, D. M., Areta, J. L., & Hawley, J. A. (2014). *Beyond muscle hypertrophy: Why dietary protein is important for endurance athletes. Applied Physiology, Nutrition, and Metabolism, 39*(9), 987–997. https://doi.org/10.1139/apnm-2013-0591

Negro, M., Giardina, S., Marzani, B., & Marzatico, F. (2008). *Branched-chain amino acid supplementation does not enhance athletic performance but affects muscle recovery and the immune system. Journal of Sports Medicine and Physical Fitness, 48*(3), 347–351. https://www.proquest.com/scholarly-journals/branched-chain-amino-acid-supplementation-does/docview/202717086/se-2

Röhling, M., McCarthy, D., & Berg, A. (2021). *Continuous protein supplementation reduces acute exercise-induced stress markers in athletes performing marathon. Nutrients*, *13*(9), 2929–. https://doi.org/10.3390/nu13092929

Vitale, K. C., & Getzin, A. (2019). *Nutrition and supplement update for the endurance athlete: Review and recommendations. Nutrients*, *11*(6), 1289–. https://doi.org/10.3390/nu11061289